



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

phyletic characters of the ferns can be estimated, the criterion of position of the nascent sorus may be held to take precedence, in point of early origin and constancy, over any soral characters except the primal features of the sporangium itself, and over any anatomical characters of the axis derivative from the protostele." This is certainly an important conclusion, and in accordance with it, the leptosporangiate ferns (exclusive of the Osmundaceae) are grouped into two series: the "Superficiales," in which the origin of the sorus is constantly from the leaf-surface; and the "Marginales," in which it is as constantly from the margin.—J. M. C.

Morphology of *Riccia*.—Miss BLACK⁴⁰ in a recent study of *Riccia Frostii* Aust. finds that this species is strictly dioicous and that the sex organs are scattered irregularly in acropetal succession. From the standpoint of the arrangement of sex organs, this indicates that *R. Frostii* is more primitive than *R. natans*, in which the antheridia are clustered in a disk, and the archegonia, which appear later, are usually in two rows. From the standpoint of restriction of antheridia and archegonia to different individuals, an advance beyond *R. natans* is clearly indicated.

Miss BLACK agrees with Miss HIRSH,⁴¹ who also studied *R. Frostii*, that the air chambers are not produced by splitting of cell walls at the angles of the cells, but by papillate outgrowth. Unfortunately, the figure given, as was the case in the work of Miss HIRSH, does not show the earliest stage in the development of the chamber, but can as easily be cited as proof that air chambers arise by splitting at the angles of the cells of the dorsal layer. The youngest air chamber shown is too old to settle the question either way, but a study of the relation of cells in the figure indicates that possibly the chamber may have arisen by splitting of the dorsal layer. This splitting need not originally occur within the tissues as some recent writers assume, but may, as DEUTSCH showed in *Targionia*, extend from the surface inward.

The rest of the investigation, which includes the development of sex organs, spermatogenesis, and sporogenesis, gives us nothing new.—W. J. G. LAND.

Peripheral leaf cells.—In many leafy liverworts there is a marked difference in form and markings of the peripheral cells of the leaf as compared with those farther away from the edge. GARJEANNE,⁴² as the result of a study of 10 genera, finds that the thickening of the peripheral cells is stronger if the plant is exposed to conditions which give great variation of water content;

⁴⁰ BLACK, CAROLINE A., The morphology of *Riccia Frostii* Aust. Ann. Botany 27:511-532. pls. 37, 38. 1913.

⁴¹ HIRSH, PAULINE E., The development of the air chambers of Ricciaceae. Bull. Torr. Bot. Club 27:73-77. figs. 6. 1910.

⁴² GARJEANNE, A. J. M., Die Randzellen einiger Jungermannienblätter. Flora 105:370-384. 1913.